

## **Child Asthma Record**

This form is to be completed by parents/carers, ideally in consultation with the child's doctor (general practitioner or specialist). Parents/carers should inform the service immediately if there are any changes to the child's asthma management. A new Asthma Record should be provided at the beginning of each year. Please tick the appropriate box, and print your answers clearly in the blank spaces where indicated.

Personal Details		
Child's Name	(first name)	(last name)
Gender □ Male □ Female	Date of Birth / / /	,
Emergency Contacts	1. Name	Relationship
(eg. Parent or Carer)	Telephone (daytime)	
	2. Name	
	Telephone (daytime)	(home)
Doctor's Contact Details	Name	Telephone
Asthma Management Plan	*	
Does the child tell the carer whe	n he/she needs medication? ☐ Yes ☐ No	
Child's Symptoms (eg cough)		
Triggers (eg exercise, pollens) _		
Medication Requirements: (Pare	nts need to supply asthma medication eg, puffer	and spacer)
Name of Medication	Method of delivery (eg puffer & spacer)	When & How Much
In an <b>EMERGENCY</b> , follow the plan that has been ticked:		
Standard Asthma First Aid Plan		My Child's Asthma First Aid Plan
First Aid Plan	remain calm and provide reassurance. one.	
First Aid Plan  Step 1: Sit the child upright and Do not leave the child alc  Step 2: Give 4 puffs of a blue rel one puff at a time, through	one. iever ( <i>Airomir, Asmol, Epaq or Ventolin</i> ), gh a spacer device*.	First Aid Plan  As written in consultation with my
First Aid Plan  Step 1: Sit the child upright and Do not leave the child alc  Step 2: Give 4 puffs of a blue rel one puff at a time, through	one. iever ( <i>Airomir, Asmol, Epaq or Ventolin</i> ),	First Aid Plan  As written in consultation with my child's doctor.  (Full details must be attached or staff will use the Standard Asthma
First Aid Plan  Step 1: Sit the child upright and Do not leave the child ale Step 2: Give 4 puffs of a blue rel one puff at a time, throug Ask the child to take 4 brows Step 3: Wait 4 minutes.  Step 4: If there is little or no improvements of the step of the	one. iever (Airomir, Asmol, Epaq or Ventolin), gh a spacer device*. reaths from the spacer after each puff. overnent, repeat steps 2 and 3. If there is ent, call an ambulance immediately (Dial 000).	First Aid Plan  As written in consultation with my child's doctor.  (Full details must be attached or staff will use the Standard Asthma
First Aid Plan  Step 1: Sit the child upright and Do not leave the child alcomposition Step 2: Give 4 puffs of a blue related one puff at a time, through Ask the child to take 4 brown Step 3: Wait 4 minutes.  Step 4: If there is little or no improvement of the steps of the steps.	one. iever (Airomir, Asmol, Epaq or Ventolin), gh a spacer device*. reaths from the spacer after each puff. ovement, repeat steps 2 and 3. If there is	First Aid Plan  As written in consultation with my child's doctor.  (Full details must be attached or staff will use the Standard Asthma
First Aid Plan  Step 1: Sit the child upright and Do not leave the child alcomposition Step 2: Give 4 puffs of a blue related one puff at a time, through Ask the child to take 4 brown Step 3: Wait 4 minutes.  Step 4: If there is little or no improvement of the steps of the steps.	one. diever (Airomir, Asmol, Epaq or Ventolin), gh a spacer device*. deaths from the spacer after each puff. deaths from the s	First Aid Plan  As written in consultation with my child's doctor.  (Full details must be attached or staff will use the Standard Asthma
Step 1: Sit the child upright and Do not leave the child ale Step 2: Give 4 puffs of a blue rel one puff at a time, throug Ask the child to take 4 brown ask the child to take 4 brown ask the child to take 4 brown ask the child to round ask the child to take 4 brown ask the ch	one. diever (Airomir, Asmol, Epaq or Ventolin), gh a spacer device*. deaths from the spacer after each puff. deaths from the s	First Aid Plan  As written in consultation with my child's doctor.  (Full details must be attached or staff will use the Standard Asthma First Aid Plan)
Step 1: Sit the child upright and Do not leave the child ale Step 2: Give 4 puffs of a blue rel one puff at a time, throug Ask the child to take 4 brown ask the child transfer ask the child to take 4 brown ask the chi	iever (Airomir, Asmol, Epaq or Ventolin), gh a spacer device*. reaths from the spacer after each puff.  ovement, repeat steps 2 and 3. If there is ent, call an ambulance immediately (Dial 000). 2 and 3 while waiting for the ambulance.  Epaq or Ventolin) on its own if no spacer is available.	As written in consultation with my child's doctor. (Full details must be attached or staff will use the Standard Asthma First Aid Plan)